

Erratum: Star formation and chemical evolution in SPH simulations: a statistical approach

Cesario Lia^{1,2}, Laura Portinari^{3,2}, and Giovanni Carraro²

¹ *SISSA/ISAS, via Beirut 4, I-34014 Trieste, Italy*

² *Dipartimento di Astronomia, Università di Padova, Vicolo dell'Osservatorio 2, I-35122 Padova, Italy*

³ *Theoretical Astrophysics Center, Juliane Maries Vej 30, DK-2100 Copenhagen Ø*

E-mail: lportina@tac.dk, carraro@pd.astro.it

Submitted: June 2002

Key words: errata, addenda — hydrodynamics — methods: numerical — stars: formation — galaxies: evolution.

The paper “Star formation and chemical evolution in SPH simulations: a statistical approach” was published on MNRAS 330, 821–836 (2002, astro-ph/0111084). Here we notify a notation error in Equation 7 and a few typos in Table 3.

Eq. 7 on p. 826 should read as follows:

$$N_{SN} \longrightarrow \frac{N_{SN}}{\int_t^{t+\Delta t} e(t') dt'}$$

with $e(t)$ given by Eq. 2 on p. 823. In fact, the particles that return to be gas within Δt represent a fraction $\int_t^{t+\Delta t} e(t') dt'$ of the whole parent stellar population, although they are assigned a return probability $g_t(\Delta t)$ (Eq. 4) to consistently account for the star particles ‘lost’ from the initial population. Only as corrected above, Eq. 7 is consistent with the corresponding metal production in Eq. 9 on p. 828.

The mistake in Eq. 7 in the paper is the result of a different layout and notation of Eqs. 3,4 between an early version of the manuscript and the final text. It is only an error of notation and in our algorithm we applied the correct expression presented here above, to obtain the correct SN rates shown e.g. in Fig. 5 and Fig. 9.

Table 3 on p. 827 contains one wrong entry (due to typesetting errors of the authors), affecting the yields of carbon $p_C(t)$ in the time range $[2 \times 10^8, 5 \times 10^9]$ yr, and a typo in the Kroupa entry for secondary nitrogen. We repeat here below the complete entries for the yields of carbon and secondary nitrogen, corrected where relevant.

chemical element	Salpeter	Kroupa	Arimoto–Yoshii	time range [yr]
¹² C $p_C(t) = \left\{ \begin{array}{l} 1.2e-10 \\ 1.2e4 t^{-2} \\ 1.1e-6 t^{-0.7} - 1.5e-13 \end{array} \right.$	1.2e-10	5.8e-11	3.6e-10	$t \in [3.4 \times 10^6, 10^7]$
	1.2e4 t^{-2}	5.8e3 t^{-2}	3.6e4 t^{-2}	$t \in [10^7, 3.4 \times 10^7]$
	1.1e-6 $t^{-0.7} - 1.5e-13$	2.9e-8 $t^{-0.5} - 4e-13$	6.4e-5 $t^{-0.9} - 1e-13$	$t \in [2 \times 10^8, 5 \times 10^9]$
¹⁴ N $p_{Ns}(t) =$	7.7 $Z_0 t^{-1.4}$	0.98 $Z_0 t^{-1.3}$	520 $Z_0 t^{-1.6}$	$t \in [3.4 \times 10^6, 15 \times 10^9]$